

OPED

Active CPM



CAMO[®]ped

The active motion device CAMO[®]ped increases the healing success rate and lessens the recovery time during therapy after surgery on the knee or hip. The CAMO[®]ped facilitates mainly passive training of the injured leg by allowing the healthy leg to compensate. Due to the so-called cross-over effect, the muscular activity of the healthy leg causes an activation of the weakened group of muscles of the injured leg. Thus, a strengthening of those muscles occurs. Further in the course of treatment,


both legs will move actively. When using CAMO[®]ped, proprioceptive activation allows for an earlier return to coordinative abilities*. The accelerated increase of the range of motion and the reduction of pain by the patient's own movement in a very early phase of recovery motivates the patient and actively supports rehabilitation.

CAMO[®]ped may also be suitable for physical thrombosis prophylaxis and as a motion device for bed-ridden patients.



*Reference: study of B. Friemert, et al. (Surgical Dept. of the Federal Armed Forces Hospital of Ulm):
"CAMO[®]ped vs. CPM – The use of motion splints in the follow-up treatment of ACL plasty"

- Foldable feet for easy transportation
- Easily adjustable angle limiter for knee joint movement between -15° hyperextension and 120° flexion
- Movement resistance regulated by 1 of 4 settings
- Telescopic reinforcement for optimal stability
- Horizontally swivel-mounted for natural knee rotation
- Adjustable resistance control for ankle movement
- Adjustment for different calf lengths



Treatment Goals

- Increased range of motion in the knee and the hip
- Increased coordination and proprioception
- Gradual strenghtening of the muscles
- Decreased swelling and effusion
- Ligament and wound healing
- Physical thrombosis prophylaxis
- Motivation of the patient through their participation in therapy

CAMO[®]ped

Indications:

- Before/after cruciate ligament plasty
- After knee and hip joint replacement
- After osteosyntheses, pseudarthroses and corrective osteosyntheses
- After reconstructive surgery on meniscus and articular cartilage
- After Synovectomy
- During prolonged confinement to bed
- During movement therapy for cardio vascular diseases and post-apoplectic
- In general for thrombosis prophylaxis

Note: CAMO[®]ped is available only by prescription!



ORDER INFORMATION

• CAMO [®] ped III, active knee mobilizer, complete	CAMO-01-3
• Set of pads, including velcro closure set, for 2 shells	POLS-02-3
• Calf shell left, complete	SCHA-04-3-LI
• Calf shell right, complete	SCHA-04-3-RE

For prices, please refer to the current price list.



For more information, please visit www.camoped.com



OPED AG

Sumpfstraße 5
CH-6312 Steinhausen
Switzerland

OPED Inc.

111 Livingston Street, Suite 1110
Brooklyn, NY 11210
USA

Fon +41(0)41/748 53 35
Fax +41(0)41/748 53 40
E-mail contact@oped.ch
www.oped.ch

Fon +1/718/210-4771
Fax +1/718/210-4751
Toll free +1/866/447-0601
email contact@opedinc.com
www.opedinc.com